

09/831656

JC18 Rec'd PCT/PTO 11 MAY 2001

Attorney Docket No.: 5753.204-US

PATENT

## IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE  
(DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371

Box PCT  
Commissioner for Patents  
Washington, DC 20231

Sir:

This is a request for filing a patent application under 35 U.S.C. 371 of the international application identified below:

INTERNATIONAL APPLICATION NO. <b>PCT/DK99/00624</b>	INTERNATIONAL FILING DATE <b>November 12, 1999</b>	PRIORITY DATE CLAIMED <b>November 12, 1998</b>
<b>APPLICANT(S)/INVENTOR(S)</b>		
Given Name (First and middle (if any))	Family Name or Surname	Residence (City and Either State or Foreign Country)
1. Jack Bech 2. Søren	1. Nielsen 2. Kjærulff	1. DK-2900 Hellerup, Denmark 2. DK-2720 Vanløse, Denmark
<b>TITLE OF THE INVENTION</b>		
Transgenic Plant Expressing Maltogenic Alpha-Amylase		

1. This is a first submission of items concerning a filing under 35 U.S.C. 371
2. This is an express request to begin national examination procedures (35 U.S.C. 371(f)).
3. The US has been elected by the expiration of 19 months from the priority date (Article 31).
4. A copy of the International Application as filed (35 U.S.C. 371(c)(2)) has been communicated by the International Bureau on June 2, 2001.
5. An oath or declaration of the inventors is submitted herewith.
6. An Assignment and Recordation Form Cover Sheet are submitted herewith.
7. A Preliminary Amendment is submitted herewith.
8. The assignee of the present application is:

Novozymes A/S  
Krogshøjvej 36  
DK-2880 Bagsvaerd  
Denmark

PCT/US/00/000000

Applicants request that the assignee information be included in the published patent application.

9. Direct all correspondence to Customer Number 25908:

PATENT TRADEMARK OFFICE



25908

10. The following fees are submitted:

BASIC NATIONAL FEE (37 CFR 1.492(a)(1)-(5)):

Search Report has been prepared by the EPO or JPO \$ 860

CLAIMS FEE

Total Claims:  $20 - 20 = 0 \times 18 =$  \$ 0

Independent Claims:  $6 - 3 = 3 \times 80 =$  \$ 240

Total Claims Fee: \$ 240

FEE FOR RECORDING ASSIGNMENT \$ 40

TOTAL NATIONAL FEE \$1,140

11. Please charge the required fee, estimated to be \$1,140, to Novozymes North America, Inc., Deposit Account No. 50-1701. A duplicate of this sheet is enclosed.

Respectfully submitted,

Date: May 11, 2001

Jason I. Garbell, Reg. No. 44,116  
Novozymes North America, Inc.  
405 Lexington Avenue, Suite 6400  
New York, NY 10174-6401  
(212) 867-0123

Attorney Docket No.: 5753.204-US

PATENT

**IN THE UNITED STATES PATENT AND TRADEMARK OFFICE**

In re Application of: Nielsen et al.

Serial No.: To Be Assigned

Confirmation No:

Group Art Unit: To Be Assigned

Filed: May 12, 2001

Examiner: To Be Assigned

For: Transgenic Plant Expressing Maltogenic Alpha-Amylase

**PRELIMINARY AMENDMENT**

Commissioner for Patents  
Washington, DC 20231

Sir:

Before the above-captioned application is taken up for examination, entry of the following amendment is respectfully requested:

**IN THE SPECIFICATION:**

At page 1, after the title, insert

**--CROSS-REFERENCE TO RELATED APPLICATIONS**

This application is a 35 U.S.C. 371 national application of PCT/DK99/00624 filed November 12, 1999 and claims, under 35 U.S.C. 119, priority of Danish application no. PA 1998 01478, filed November 12, 1998, and the benefit of U.S. application no. 60/123,643, filed March 10, 1999, the contents of which are fully incorporated herein by reference.--

**IN THE CLAIMS:**

Please cancel claims 1-34 without prejudice or disclaimer.

Please add the following new claims:

--35. A transgenic cereal plant cell comprising a nucleotide sequence encoding a maltogenic alpha-amylase.--

--36. The plant cell according to claim 35, wherein the plant cell is a wheat plant cell.--

RECEIVED  
MAY 15 2001  
PCT/PTO

--37. The plant cell according to claim 35, wherein the maltogenic alpha-amylase has the amino acid sequence shown in SEQ ID NO: 2 or the amino sequence acid sequence of amino acids 1-686 of SEQ ID NO:1.--

--38. The plant cell according to claim 35, wherein the maltogenic alpha-amylase has an amino acid sequence which has:

- i) at least 70% identity to SEQ ID NO: 2; or
- ii) at least 70% identity to the amino acid sequence set forth in amino acids 1-686 of SEQ ID NO:1.--

--39. The plant cell according to claim 35, wherein said wherein the nucleotide sequence is operably linked to a seed specific promoter.--

--40. The plant cell according to claim 35, wherein the nucleotide sequence encoding the maltogenic alpha-amylase is derived from a microorganism.--

--41. The plant cell according to claim 40, wherein the nucleotide sequence encoding the maltogenic alpha-amylase is derived from the *Bacillus* strain NCIB 11837.--

--42. A transgenic cereal plant regenerated from a plant cell of claim 35 and the progeny of the plant, wherein the plant and the progeny of the plant are capable of expressing maltogenic alpha-amylase in the seeds of the plant or the progeny of the plant.--

--43. A transgenic cereal plant comprising a nucleotide sequence encoding a maltogenic alpha-amylase. --

--44. The plant according to claim 43 which is a wheat plant.--

--45. The plant according to claim 43, wherein the maltogenic amylase is a maltogenic alpha-amylase having:

- (a) the amino acid sequence shown in SEQ ID NO: 2;
- (b) the amino sequence acid sequence of amino acids 1-686 of SEQ ID NO:1;
- (c) an amino acid sequence which has at least 70% identity to SEQ ID NO: 2; or
- (d) an amino acid sequence which has at least at least 70% identity to the amino acid sequence set forth in amino acids 1-686 of SEQ ID NO:1.--

--46. A seed of the cereal plant of claim 43, wherein the seed includes maltogenic alpha-amylase in an amount effective to delay staling of bread baked from the seed.--

--47. A transgenic cereal seed comprising a maltogenic alpha-amylase in an amount effective to delay staling of bread baked from the seed.--

--48. The seed of claim 46, wherein the maltogenic alpha-amylase is a maltogenic alpha-amylase having:

- (a) the amino acid sequence shown in SEQ ID NO: 2;
- (b) the amino sequence acid sequence of amino acids 1-686 of SEQ ID NO:1;
- (c) an amino acid sequence which has at least 70% identity to SEQ ID NO: 2; or
- (d) an amino acid sequence which has at least at least 70% identity to the amino acid sequence set forth in amino acids 1-686 of SEQ ID NO:1.--

--49. The seed of claim 46, wherein the seed is a wheat seed.--

--50. A method for preparing a baked product, comprising the steps of:

- i) expressing a maltogenic alpha-amylase in the seed of a transgenic cereal plant;
- ii) preparing flour from said seed comprising said maltogenic alpha-amylase;
- iii) preparing a dough comprising the flour of step ii); and
- iv) baking the dough to obtain a baked product.--

--51. A method for preparing a baked product, comprising the steps of:

- i) preparing flour from cereal seed, said seed comprising a maltogenic alpha-amylase;
- ii) preparing a dough comprising the flour of step i); and
- iii) baking the dough to obtain a baked product.--

--52. A method for preparing a baked product, comprising the steps of:

- i) preparing a dough from flour obtained from cereal seed, said seed comprising a maltogenic alpha-amylase;
- ii) preparing a dough comprising the flour of step i); and
- iii) baking the dough to obtain a baked product.--

--53. The method according of claim 50, wherein the maltogenic alpha-amylase is a maltogenic alpha-amylase having:

- (e) the amino acid sequence shown in SEQ ID NO: 2;

- (f) the amino sequence acid sequence of amino acids 1-686 of SEQ ID NO:1;
- (g) an amino acid sequence which has at least 70% identity to SEQ ID NO: 2; or
- (h) an amino acid sequence which has at least at least 70% identity to the amino acid sequence set forth in amino acids 1-686 of SEQ ID NO:1.--

--54. The method according to claim 50, wherein the seed includes the maltogenic alpha-amylase in an amount effective to delay staling of the bread product.--


#### REMARKS

Claims 1-34 are canceled. New claims 35-54 are added. New claims 35-54 are supported by the specification and claims as originally filed. No new matter has been added and entry of the amendment is respectfully requested.

The Examiner is hereby invited to contact the undersigned by telephone if there are any questions concerning this amendment or application.

Respectfully submitted,

Date: May 11, 2001

  
\_\_\_\_\_  
Jason I. Garbell, Reg. No. 44,116  
Novozymes North America, Inc.  
405 Lexington Avenue, Suite 6400  
New York, NY 10174-6401  
(212) 867-0123

## TRANSGENIC PLANT EXPRESSING MALTOGENIC ALPHA-AMYLASE

## FIELD OF THE INVENTION

5 The present invention relates to a transgenic plant cell expressing a maltogenic amylase or a beta-amylase, a transgenic plant regenerated from said cell, seeds comprising a maltogenic amylase or a beta-amylase and the use of said seeds, optionally in ground form, for catalyzing an industrial process.

10

## BACKGROUND OF THE INVENTION

Maltogenic alpha-amylase (glucan 1,4- $\alpha$ -maltohydrolase, E.C. 3.2.1.133) is able to hydrolyze amylose and amylopectin to maltose in the alpha-configuration, and is also able to  
15 hydrolyze maltotriose as well as cyclodextrin. A maltogenic alpha-amylase from *Bacillus* (EP 120 693) is commercially available under the trade name Novamyl<sup>®</sup> (product of Novo Nordisk A/S, Denmark) and is widely used in the baking industry as an anti-staling agent due to its ability to reduce  
20 retrogradation of starch/amylopectin. Novamyl is further described by Christophersen, C., et al., 1998, Starch 50, pp 39-45. Variants of Novamyl<sup>®</sup> and the three-dimensional structure of Novamyl<sup>®</sup> are disclosed in WO 99/43794.

25 WO 91/14772 discloses transgenic seeds expressing enhanced amounts of enzymes, and the use of such seeds in catalyzing industrial processes. Baking is mentioned as one example of an industrial process for which  $\alpha$ -amylase can be used and it is stated that the seeds may be ground before being incorporated  
30 into flour.

Vickers et al, Journal of the Institute of Brewing, Vol. 102, No. 2 pp. 75-78 (1996) speculate in using a *Bacillus licheniformis*  $\alpha$ -amylase as a candidate enzyme for the genetic  
35 transformation of malting barley.

## BRIEF DESCRIPTION OF THE INVENTION

The present invention relates to a cell of a transgenic seed producing plant transformed with at least one nucleotide  
5 sequence encoding a maltogenic alpha-amylase or a beta-amylase which, in the cell, is operably linked to elements required for mediating expression from said nucleotide sequence in the seeds of a plant regenerated from the plant cell.

10 In a further aspect the invention relates to a transgenic seed-producing plant regenerated from a cell of the invention and expressing measurable quantities of a maltogenic alpha-amylase or a beta-amylase in its seeds.

15 In a still further aspect the invention relates to the seeds of a plant of the invention, optionally in ground form, and the use of such seed for catalyzing an industrial process.

The invention also relates to a method for producing a  
20 maltogenic alpha-amylase or beta-amylase comprising recovering the amylase or the beta-amylase from seeds of the invention.

## DETAILED DISCLOSURE OF THE INVENTION

25 The maltogenic alpha-amylase is an enzyme classified in EC 3.2.1.133. The enzymatic activity does not require a non-reducing end on the substrate and the primary enzymatic activity results in the degradation of amylopectin and amylose to maltose and longer maltodextrins. It is able to hydrolyze  
30 amylose and amylopectin to maltose in the alpha-configuration, and is also able to hydrolyze maltotriose as well as cyclodextrin.

For the present invention in a particularly interesting  
35 embodiment the maltogenic alpha-amylase enzyme corresponds to maltogenic alpha-amylase cloned from Bacillus as described in EP 120 693 (hereinafter referred to as Novamyl). Novamyl has the amino acid sequence set forth in amino acids 1-686 of SEQ



ID NO: 1. Novamyl is encoded in the gene harboured in the Bacillus strain NCIB 11837 (c.f. EP 120 693) which has the nucleic acid sequence set forth in SEQ ID NO:1. Thus, in one preferred embodiment of the invention the maltogenic alpha-amylase enzyme is identical to a maltogenic alpha-amylase obtainable from Bacillus strain NCIB 11837. In the context of the invention is also contemplated a nucleotide sequence encoding said enzyme, such as e.g. a nucleotide sequence obtainable from Bacillus strain NCIB 11837 encoding said enzyme. The coding sequence for Novamyl may be obtained from the strain DSM 11837 or from the plasmid denoted pLBei010 as indicated in WO 99/43794. The plasmid pLBei010 contains amyM in which the expression of amyM is directed by its own promoter and the complete gene encoding Novamyl, e.g., as contained in the strain DSM 11837. The plasmid contains the origin of replication, ori, from plasmid pUB110 and an kanamycin resistance marker for selection purposes. pLBei010 is shown in Fig. 1. Preferably the maltogenic alpha-amylase enzyme for the present invention has an anti-staling effect in baking.

The present inventors have found that maltogenic alpha-amylases, (such as, e.g. Novamyl®) has a very unique performance in bread making. Other thermostable  $\alpha$ -amylases like BAN® (product of Novo Nordisk A/S) or Termamyl® (product of Novo Nordisk A/S) must be dosed very carefully in tight intervals, e.g. between 0.5-2 times of the optimum dosage in a given recipe. Otherwise the risk is high that there is either no effect (low dosage) or too high effect (high dosage). The latter results in a gummy, non-elastic and sticky crumb, unsuited for eating. The inventors have found that maltogenic alpha-amylases does not have this problem, but can be dosed broadly. For instance, Novamyl® has a positive function on e.g. staling properties from a level of e.g. 200 MANU/kg flour to 5.000 MANU/kg, i.e. a much safer amylase in practical application than other  $\alpha$ -amylases. This property of Novamyl® is thus a fundamental difference compared to known  $\alpha$ -amylases. This characteristic makes it superior in baking applications, such as in connection with anti-staling, and it is also

contemplated that this property makes maltogenic alpha--  
amylases, such as e.g. Novamyl®, particularly suitable for  
transgenic expression in plants, in particular in a seed  
producing plant, such as e.g. wheat. Accordingly, one  
5 embodiment of the present invention relates to the expression  
of a maltogenic alpha-amylase, such as Novamyl, in the seeds of  
a seed producing plant, such as, e.g. wheat.

Of particular interest for the present invention is enzymes  
10 having an anti-staling effect and at the same time having the  
above indicated characteristic, i.e. that relatively high  
dosage of the enzyme in baking does not have an essentially  
negative effect as compared to other enzymes. Within the scope  
of the invention is expression of such enzymes (with anti-  
15 staling effect and the above indicated dosage characteristics  
in baking) in the seeds of a seed producing plant, such as,  
e.g. wheat, and the use of such seed in baking. Examples of  
enzymes with such characteristics are maltogenic alpha-  
amylases, such as Novamyl. The negative effect of high dosage  
20 may be exemplified by the effect of use of high dosage of  $\alpha$ -  
amylases in baking as indicated above.

Beta-amylase is another example of an enzyme that shows a  
relatively low level of criticality to high dosage in baking.  
25 Thus, one embodiment of the present invention relates to the  
expression of a beta-amylase in the seeds of a seed producing  
plant, such as, e.g. wheat, and the use of such seed in baking.

Within the scope of the invention is a maltogenic alpha-amylase  
30 being this an enzyme with one or more characteristics selected  
from the group consisting of:

i) having the amino acid sequence set forth in SEQ ID NO:2;

35 ii) having the amino acid sequence set forth in amino acids 1-  
686 of SEQ ID NO:1;

iii) having a three dimensional structural homology to Novamyl;

iv) having an amino acid sequence which has at least 70% identity to SEQ ID NO: 2, preferably at least 75%, 80 %, 85% or at least 90%, e.g. at least 95%, 97%, 98 %, or at least 99%;

v) having an amino acid sequence which has at least 70% identity to the amino acid sequence set forth in amino acids 1-686 of SEQ ID NO:1, preferably at least 75%, 80 %, 85% or at least 90%, e.g. at least 95%, 97%, 98 %, or at least 99%;

vi) a fragment of i), ii) iv) or v), said fragment consisting of 10-600 amino acid residues, such as in the range of 30-300 amino acid residues, such as 50-100 amino acid residues;

vii) an amino acid sequence encoded by a nucleotide sequence which hybridizes (1) to the DNA sequence set forth in SEQ ID NO:1, (2) to the DNA sequence encoding Novamyl harboured in the *Bacillus* strain NCIB 11837, (3) to the DNA sequence contained in the nucleotides 100 to 2157 of SEQ ID NO:1, (4) to a subsequence of (1) or (3) of at least 30 nucleotides, such as at least 50 nucleotides, at least 100 nucleotides, at least 200 nucleotides, or (5) to a complementary strand of (1), (3), or (4) under low stringency conditions, or under medium stringency, more preferably at medium/high stringency or at high stringency or even more preferably at very high stringency;

viii) a catalytic binding site comprising amino acid residues similar to D229, E257 and D328 as shown in the amino acid sequence set forth in amino acids 1-686 of SEQ ID NO:1;

ix) a variant of the amino acid sequence set forth in amino acids 1-686 of SEQ ID NO:1 comprising a substitution, deletion, and/or insertion of one or more amino acids;

The structural homology referred to above in iii) is as disclosed in WO 99/43794 and is based on other sequence homologies, hydrophobic cluster analysis or by reverse threading

(Huber, T ; Torda, AE, PROTEIN SCIENCE Vol. 7 , No. 1 pp. 142-149 (1998)) and which by any of these methods is predicted to have the same tertiary structure as Novamyl, wherein the tertiary structure refers to the overall folding or the folding of Domains A, B, and C, more preferably including Domain D, and most preferably including Domain E as disclosed in WO 99/43794. Alternatively, a structural alignment between Novamyl and a maltogenic alpha-amylase may be used to identify equivalent positions.

Maltogenic alpha-amylase variants are described in WO 99/43794. In further embodiments of the present invention the maltogenic alpha-amylase enzyme is a variant of the amino acid sequence set forth in amino acids 1-686 of SEQ ID NO:1 or a variant of SEQ ID NO:2, such variants are disclosed in WO 99/43794. WO 99/43794 also discloses how such suitable modifications may be identified and how to prepare the modifications. Accordingly, the maltogenic alpha-amylase enzyme of the invention may be a maltogenic alpha-amylase enzyme variant having a modified amino acid sequence compared the amino acid sequence set forth in amino acids 1-686 of SEQ ID NO:1 or compared to SEQ ID NO:2.

The maltogenic alpha-amylase enzyme variant may have one or more of the following properties which are modified compared to an enzyme having the amino acid sequence of set forth in amino acids 1-686 of SEQ ID NO:1 or compared to SEQ ID NO: 2, such as stability (e.g. thermostability), pH dependent activity, substrate specificity, specific activity or ability to reduce retrogradation of starch or staling of bread. Thus, the altered property may be an altered specific activity at a given pH and/or an altered substrate specificity, such as an altered pattern of substrate cleavage or an altered pattern of substrate inhibition.

These variants may be modifications of SEQ ID NO: 2 or the amino acid sequence set forth in amino acids 1-686 of SEQ ID NO:1 consisting in substitution, deletion or insertion, or a mixture of these, of one or more amino acid residues.

In further embodiments of the invention the variant of a maltogenic alpha-amylase has an altered pH dependent activity profile as compared to Novamyl and has an amino acid sequence comprising a modification of an amino acid residue corresponding to one or more of the following residues of the amino acid sequence set forth in SEQ ID NO: 1:

D127, V129, F188, A229, Y258, V281, F284, T288, N327, M330, G370, N371, and D372, L71, S72, V74, L75, L78, T80, L81, G83, T84, D85, N86, T87, G88, Y89, H90, G91, T94, R95, D96, F97, I174, S175, N176, D178, D179, R180, Y181, E182, A183, Q184, K186, N187, F188, T189, D190, A192, G193, F194, S195, L196.

In further embodiments of the present invention the variant comprises a modification corresponding to one or more of the following modifications in the amino acid sequence set forth in SEQ ID NO: 1: D127N/L, V129S/T/G/V, F188E/K/H, A229S/T/G/V, Y258E/D/K/R/F/N, V281L/T, F284K/H/D/E/Y, T288E/K/R, N327D, M330L/F/I/D/E/K, G370N, N371D/E/G/K, and D372N/V, L71I, S72C, V74I, L75N/D/Q/I/V, L78N/I, T80I/L/V/S/N/G, L81I/V/S/T/N/Q/K/H, G83A/S/T/N/Q/E/D/R/H/L, T84S/A/N/D/G, D85A/T/S/N/G, N86Q/E/D/Y/H/K, T87S/I, G88A/S/T, Y89F, H90N/Q/K, G91A/S/T, T94N/D/A/M/V/I, R95K/Q, D96N/V/Q/I, F97Y, I174N/Q/L, S175T/A/N/D, N176S/T/H/Q/P, D178N/Q/E/K/H, D179Y/N/H, R180W, Y181R/F/C/L, E182D, A183S/C/G, Q184E, K186R, N187Q/E/L/F/H/K/V/L, F188Y/L/I/H/N, T189N/D/A/S/H/Y/G, D190E/Q/H/N/K, A192T/D/E/N/K, G193A/S/T, F194Y, S195N/D/E/R/K/G, L196I.

Other variants contemplated for the present invention are variants of Novamyl having an altered  $\text{Ca}^{2+}$  binding as compared to the parent maltogenic alpha-amylase and where said variant has an amino acid sequence comprising a modification of an amino acid residue corresponding to one or more of the following residues of the amino acid sequence set forth in SEQ ID NO: 1: D17, A30, S32, R95, H103, N131, Q201, I174, and/or H169, V74, L75, L78, T80, L81, T87, G88, Y89, H90, G91, T94, R95, D96, F97, Y167, F168, H169, H170, N171, G172, D173, I174, S175, N176,

D178, D179, R180, Y181, E182, A183, Q184, K186, N187, F188, T189.

In one embodiment, the variant of SEQ ID NO: 1 has an altered  
 5 Ca<sup>2+</sup> binding as compared to the parent maltogenic alpha-amylase. In one embodiment of the invention the Ca<sup>2+</sup> binding of a maltogenic alpha-amylase is change the partial sequence N28-P29-A30-K31-S32-Y33-G34 as set forth in SEQ ID NO: 1 is modified.

10

Further contemplated variants are variants having an amino acid sequence comprising a substitution corresponding to one or more of the following substitutions in the amino acid sequence set forth in SEQ ID NO: 1:

15 D17E/Q, A30M/L/A/V/I/E/Q, S32D/E/N/Q, R95M/L/A/V/I/E/Q, H103Y/N/Q/D/E, N131D, Q201E, I174E/Q, and H169N/D/E/Q, V74I, L75N/D/Q/I/V, L78N/I, T80I/L/V/S/N/G, L81I/V/S/T/N/Q/K/H, T87S/I, G88A/S/T, Y89F, H90N/Q/K, G91A/S/T, T94N/D/A/M/V/I, R95K/Q, D96N/V/Q/I, F97Y, Y167F/R/C, F168Y, H169N/Q/K,  
 20 H170N/Q/K, N171D/E/Q/H/R/K/G, G172A/T/S, D173N/S/T/Y/R/G, I174N/Q/L, S175T/A/N/D, N176S/T/H/Q/P, D178N/Q/E/K/H, D179Y/N/H, R180W, Y181R/F/C/L, E182D, A183S/C/G, Q184E, K186R, N187Q/E/L/F/H/K/V/L, F188Y/L/I/H/N, T189N/D/A/S/H/Y/G.

25 The maltogenic alpha-amylase variants may also have an altered thermostability and/or an altered temperature dependent activity profile compared to Novamyl. Such variants may comprise a substitution of an amino acid residue corresponding to one or more of the following residues of the amino acid sequence set  
 30 forth in SEQ ID NO: 1:

L51, L75, L78, G88, G91, T94, V114, I125, V126, T134, G157, L217, S235, G236, V254, V279, V281, L286, V289, I290, V308, L321, I325, D326, L343, F349, S353, I359, I405, L448, Q449, L452, I470, G509, V515, S583, G625, L627, L628 and A670, L71,  
 35 S72, V74, L75, L78, T80, L81, G83, T84, D85, N86, T87, G88, Y89, H90, G91, T94, R95, D96, F97, Y167, F168, H169, H170, N171, G172, D173, I174, S175, N176, D178, D179, R180, Y181, E182,

A183, Q184, K186, N187, F188, T189, D190, A192, G193, F194, S195, L196.

Such variants in the context of the invention may comprise one  
 5 or more substitutions corresponding to the following  
 substitutions in the amino acid sequence set forth in SEQ ID NO:  
 1: L217 in combination with L75 (e.g. L217F/Y in combination  
 with L75F/Y), L51W, L75F/Y, L78I, G88A/V/T, G91T/S/V/N,  
 T94V/I/L, V114V/I/L, I125L/M/F/Y/W, V126I/L, T134V/I/L/M/F/Y/W,  
 10 G157A/V/I/L, L217V/I/M/F/Y/W, S235I/L/M/F/Y/W,  
 G236A/V/I/L/M/F/Y/W, V254I/L/M/F/Y/W, V279M/I/L/F,  
 V281I/L/M/F/Y/W, L286F, V289I/L/R, I290M/L/F, V308I/L/M/F/Y/W,  
 L321I/M/F/Y/W, I325L/M/F/Y/W, D326E/Q, L343M/F/Y/W, F349W/Y,  
 S353V/I/L, I359L/M/F/Y/W, I405M/L/Y/F/W, L448Y, Q449Y,  
 15 L452M/Y/F/W, I470M/L/F, G509A/V/I/L/M/S/T/D/N, V515I/L,  
 S583V/I/L/V, G625A/V/I/L/M/F/Y/W, L627M/F/Y, L628M/I/F/Y/W and  
 A670V/I/L/M/F/Y/W, L71I, S72C, V74I, L75N/D/Q/I/V, L78N/I,  
 T80I/L/V/S/N/G, L81I/V/S/T/N/Q/K/H, G83A/S/T/N/Q/E/D/R/H/L,  
 T84S/A/N/D/G, D85A/T/S/N/G, N86Q/E/D/Y/H/K, T87S/I, G88A/S/T,  
 20 Y89F, H90N/Q/K, G91A/S/T, T94N/D/A/M/V/I, R95K/Q, D96N/V/Q/I,  
 F97Y, Y167F/R/C, F168Y, H169N/Q/K, H170N/Q/K, N171D/E/Q/H/R/K/G,  
 G172A/T/S, D173N/S/T/Y/R/G, I174N/Q/L, S175T/A/N/D,  
 N176S/T/H/Q/P, D178N/Q/E/K/H, D179Y/N/H, R180W, Y181R/F/C/L,  
 E182D, A183S/C/G, Q184E, K186R, N187Q/E/L/F/H/K/V/L,  
 25 F188Y/L/I/H/N, T189N/D/A/S/H/Y/G, D190E/Q/H/N/K, A192T/D/E/N/K,  
 G193A/S/T, F194Y, S195N/D/E/R/K/G, L196I.

Further variants may have an altered substrate binding site as  
 compared to said parent. Such variant may comprise a  
 30 modification in a position corresponding to one or both of the  
 following positions in SEQ ID NO: 1: V281 and/or A629. In one  
 embodiment of the invention the variant comprises a modification  
 corresponding to: V281Q and/or A629N/D/E/Q.

35 Maltogenic alpha amylases having an improved ability to reduce  
 the retrogradation of starch and/or the staling of bread  
 compared to Novamyl is also contemplated within the context of  
 the invention. Such variants of Novamyl may comprise a

modification at one or more positions corresponding to the following amino acid residues in SEQ ID NO: 1:

A30, K40, N115, T142, F188, T189, P191, A192, G193, F194, S195, D261, N327, K425, K520 and N595. Maltogenic alpha-amylases in the context of the invention may be a variant of Novamyl comprising one or more modifications corresponding to the following in SEQ ID NO: 1: A30D, K40R, N115D, T142A, F188L, T189Y, Δ (191-195), D261G, D261G, N327S, K425E, K520R and N595I.

Within the context of the invention are variants having a combination of one or more of the above with any of the other modifications disclosed herein.

Thus, the maltogenic alpha-amylase in relation to the present invention may have an amino acid which is modified compared to Novamyl where said modified sequence has one or more of the following modifications compared to the amino acid sequence set forth in amino acids 1-686 of SEQ ID NO:1:

192-A-193; Δ (191-195); D17E; S32Q; S32D; S32N; H103Y; N131D; I174Q; I174E; N176S; F188H; F188E; Δ 191; 192-A-193; 192-A-G-193; Δ 192; Δ 262-266; F284E; F284D; F284K; T288K; T288R; N327D; G397P;

N115D+ F188L; T142A+ D261G; G370N+ N371G; N115D+ F188L; A30D+ K40R+ D261G; F188L+ V336L+ T525A; F188I+ Y422F+ I660V; F188L+ D261G+ T288P; Δ (191-195)+ F188L+ T189Y; K40R+ F188L+ D261G+ A483T; T142A+ N327S+ K425E+ K520R+ N595I; T142A+ N327S+ K425E+ K520R+ N595I.

*Nomenclature for amino acid modifications:* The nomenclature used herein for defining mutations is essentially as described in WO 92/05249. Thus, F188H indicates a substitution of the amino acid F (Phe) in position 188 with the amino acid H (His). V129S/T/G/V indicates a substitution of V129 with S, T, G or V. Δ (191-195) or Δ (191-195) indicates a deletion of amino acids in positions 191-195. 192-A-193 indicates an insertion of A between amino acids 192 and 193.

The polypeptide sequence identity referred to above in v) is determined as the degree of homology between two sequences



indicating a derivation of the first sequence from the second. The identity may be suitably determined by means of computer programs known in the art such as GAP, provided in the GCG program package (Program Manual for the Wisconsin Package, Version 8, August 1994, Genetics Computer Group, 575 Science Drive, Madison, Wisconsin, USA 53711; Needleman, S.B. and Wunsch, C.D., 1970, Journal of Molecular Biology, 48, 443-453). Using GAP with the following settings for polypeptide sequence comparison: GAP creation penalty of 3.0 and GAP extension penalty of 0.1, the mature protein part of a polypeptide encoded by an analogous DNA sequence of the invention exhibits a degree of identity preferably of at least 40%, preferably at least 50%, least 60%, at least 67%, at least 70%, , preferably at least 75%, 80 %, 85%, at least 90%, e.g. at least 95%, 97%, 98 %, or at least 99% identity to the amino acid sequence set forth in amino acids 1-686 of SEQ ID NO:1 or to the amino acid sequence set forth in SEQ ID NO:2. In a preferred embodiment of the invention, the degree of identity between two amino acid sequences as disclosed herein is determined by the Clustal method (Higgins, 1989, CABIOS 5: 151-153) using the LASERGENE™ MEGALIGN™ software (DNASTAR, Inc., Madison, WI) with an identity table and the following multiple alignment parameters: Gap penalty of 10 and gap length penalty of 10. Pairwise alignment parameters were Ktuple=1, gap penalty=3, windows=5, and diagonals=5].

In connection with maltogenic alpha-amylases characterised by vii), the oligonucleotide probe used in a hybridization may be suitably prepared on the basis of the nucleic acid sequence set forth in SEQ ID NO:1.

The hybridization referred to above in vii) is intended to indicate that the analogous DNA sequence hybridizes to the nucleotide probe corresponding to the protein encoding part of the nucleic sequence shown in SEQ ID NO:1, under at least low stringency conditions as described in detail below.

Suitable experimental conditions for determining hybridization at low stringency between a nucleotide probe and a homologous DNA or RNA sequence involves presoaking of the filter containing the DNA fragments or RNA to hybridize in 5x SSC (sodium chloride/sodium citrate, Sambrook, et al., 1989) for 10 min, and prehybridization of the filter in a solution of 5x SSC, 5x Denhardt's solution (Sambrook, et al., 1989), 0.5% SDS and 100 µg/ml of denatured sonicated salmon sperm DNA (Sambrook, et al., 1989), followed by hybridization in the same solution containing a random-primed (Feinberg, A. P. and Vogelstein, B. (1983) *Anal. Biochem.* 132:6-13), <sup>32</sup>P-dCTP-labeled (specific activity > 1 x 10<sup>9</sup> cpm/µg) probe for 12 hours at ca. 45°C. The filter is then washed twice for 30 minutes in 2x SSC, 0.5% SDS at least 55°C (low stringency), more preferably at least 60°C (medium stringency), more preferably at least 65°C (medium/high stringency), more preferably at least 70°C (high stringency), even more preferably at least 75°C (very high stringency). Molecules which hybridize to the oligonucleotide probe under these conditions are detected by exposure to x-ray film.

The following paragraphs describes how to prepare the transgenic plants of the invention, i.e. plants transformed so as to produce the enzymes as disclosed herein. Mainly maltogenic alpha-amylase is use as an examples but it is considered to be equally valid for the other enzymes mentioned herein, such as e.g. a beta-amylase.

### 30 *Cloning a DNA sequence encoding a maltogenic alpha-amylase*

The nucleotide sequence encoding the enzyme of the invention, such as the maltogenic alpha-amylase or beta-amylase, may be of any origin, including mammalian, plant and microbial origin and may be isolated from these sources by conventional methods.

35 Preferably, the nucleotide sequence is derived from a microorganism, such as a fungus, e.g. a yeast or a filamentous fungus, or a bacterium. The DNA sequence encoding a parent

0931655-090201

maltogenic alpha-amylase may be isolated from the cell producing the maltogenic alpha-amylase in question, using various methods well known in the art, for example, from the *Bacillus* strain NCIB 11837. First, a genomic DNA and/or cDNA library should be constructed using chromosomal DNA or messenger RNA from the organism that produces the maltogenic alpha-amylase to be studied. Then, if the amino acid sequence of the maltogenic alpha-amylase is known, homologous, labelled oligonucleotide probes may be synthesised and used to identify maltogenic alpha-amylase-encoding clones from a genomic library prepared from the organism in question. Alternatively, a labelled oligonucleotide probe containing sequences homologous to a known maltogenic alpha-amylase gene could be used as a probe to identify maltogenic alpha amylase-encoding clones, using hybridization and washing conditions of lower stringency.

Another method for identifying maltogenic alpha amylase-encoding clones involves inserting fragments of genomic DNA into an expression vector, such as a plasmid, transforming maltogenic alpha-amylase negative bacteria with the resulting genomic DNA library, and then plating the transformed bacteria onto agar containing a substrate for maltogenic alpha-amylase, thereby allowing clones expressing maltogenic alpha-amylase activity to be identified.

Alternatively, the DNA sequence encoding the enzyme may be prepared synthetically by established standard methods, e.g. the phosphoroamidite method described by S.L. Beaucage and M.H. Caruthers (1981) or the method described by Matthes et al. (1984). In the phosphoroamidite method, oligonucleotides are synthesized, e.g. in an automatic DNA synthesizer, purified, annealed, ligated and cloned in appropriate vectors.

Finally, the DNA sequence may be of mixed genomic and synthetic origin, mixed synthetic and cDNA origin or mixed genomic and cDNA origin, prepared by ligating fragments of synthetic, genomic or cDNA origin, wherein the fragments correspond to various parts of the entire DNA sequence, in accordance with

techniques well known in the art. The DNA sequence may also be prepared by polymerase chain reaction (PCR) using specific primers, for instance as described in US 4,683,202 or R.K. Saiki et al. (1988). See also WO 99/43794 disclosing how to make  
5 variants, e.g. by use of mutagenesis techniques known in the art.

#### *Expression Constructs*

In order to accomplish expression of the maltogenic alpha-amylase in seeds of the transgenic plant of the invention the  
10 nucleotide sequence encoding the amylase is inserted into an expression construct containing regulatory elements capable of directing the expression of the nucleotide sequence and, if necessary, to direct secretion of the gene product or targeting  
15 of the gene product to the seeds of the plant. Manipulation of nucleotide sequences using restriction endonucleases to cleave DNA molecules into fragments and DNA ligase enzymes to unite compatible fragments into a single DNA molecule with subsequent incorporation into a suitable plasmid, cosmid, or other  
20 transformation vector are well-known in the art.

In order for transcription to occur the nucleotide sequence encoding the maltogenic alpha-amylase is operably linked to a suitable promoter capable of mediating transcription in the  
25 plant in question. The promoter may be an inducible promoter or a constitutive promoter. Typically, an inducible promoter mediates transcription in a tissue-specific or growth-stage specific manner, whereas a constitutive promoter provides for sustained transcription in all cell tissues. An example of a  
30 suitable constitutive promoter useful for the present invention is the cauliflower mosaic virus 35 S promoter. Other constitutive promoters are transcription initiation sequences from the tumor-inducing plasmid (Ti) of *Agrobacterium* such as the octopine synthase, nopaline synthase, or mannopine  
35 synthase initiator.

Examples of suitable inducible promoters include a seed-specific promoter such as the promoter expressing  $\alpha$ -amylase in

wheat seeds (see Stefanov et al, Acta Biologica Hungarica Vol. 42, No. 4 pp. 323-330 (1991)), a promoter of the gene encoding a rice seed storage protein such as glutelin, prolamin, globulin or albumin (Wu et al., Plant and Cell Physiology Vol. 39, No. 8 pp. 885-889 (1998)), a *Vicia faba* promoter from the legumin B4 and the unknown seed protein gene from *Vicia faba* described by Conrad U. et al, Journal of Plant Physiology Vol. 152, No. 6 pp. 708-711 (1998), the storage protein napA promoter from Brassica napus, or any other seed specific promoter known in the art, e.g. as described in WO 91/14772.

In order to increase the expression of the maltogenic alpha-amylase it is desirable that a promoter enhancer element is used. For instance, the promoter enhancer may be an intron which is placed between the promoter and the amylase gene. The intron may be one derived from a monocot or a dicot. For instance, the intron may be the first intron from the rice Waxy (Wx) gene (Li et al., Plant Science Vol. 108, No. 2, pp. 181-190 (1995)), the first intron from the maize Ubi1 (Ubiquitin) gene (Vain et al., Plant Cell Reports Vol. 15, No. 7 pp. 489-494 (1996)) or the first intron from the Act1 (actin) gene. As an example of a dicot intron the chsA intron (Vain et al. op cit.) is mentioned. Also, a seed specific enhancer may be used to increase the expression of the maltogenic alpha-amylase in seeds. An example of a seed specific enhancer is the one derived from the beta-phaseolin gene encoding the major seed storage protein of bean (*Phaseolus vulgaris*) disclosed by Vandergeest and Hall, Plant Molecular Biology Vol. 32, No. 4, pp. 579-588 (1996).

Also, the expression construct contains a terminator sequence to signal transcription termination of the maltogenic alpha-amylase gene such as the rbcS2' and the nos3' terminators.

To facilitate selection of successfully transformed plants, the expression construct should also include one or more selectable markers, e.g. an antibiotic resistance selection marker or a selection marker providing resistance to a herbicide. One

widely used selection marker is the neomycin phosphotransferase gene (NPTII) which provides kanamycin resistance. Examples of other suitable markers include a marker providing a measurable enzyme activity, e.g. dihydrofolate reductase, luciferase, and  $\beta$ -glucoronidase (GUS). Phosphinothricin acetyl transferase may be used as a selection marker in combination with the herbicide basta or bialaphos.

#### *Transgenic plant species*

In the present context the term "transgenic plant" is intended to mean a plant which has been genetically modified to express a maltogenic alpha-amylase and progeny of such plant having retained the capability of producing a maltogenic alpha-amylase. The term also includes a part of such plant such as a leaf, seed, stem, any tissue from the plant, an organelle, a cell of the plant, etc.

Any transformable seed-producing plant species may be used for the present invention. Of particular interest is a monocotyledonous plant species, in particular crop or cereal plants such as wheat (*Triticum*, e.g. *aestivum*), barley (*Hordeum*, e.g. *vulgare*), oats, rye, rice, sorghum and corn (*Zea*, eg *mays*). In particular, wheat is preferred.

#### *Transformation of plants*

The transgenic plant cell of the invention may be prepared by methods known in the art. The transformation method used will depend on the plant species to be transformed and can be selected from any of the transformation methods known in the art such as *Agrobacterium* mediated transformation (Zambryski et al., EMBO Journal 2, pp 2143-2150, 1993), particle bombardment (Vasil et al. 1991), electroporation (Fromm et al. 1986, Nature 319, pp 791-793), and virus mediated transformation. For transformation of monocots particle bombardment (i.e. biolistic transformation) of embryogenic cell lines or cultured embryos are preferred. In the following references disclosing methods for transforming different plants are mentioned together with the plant: Rice (Cristou et al.

1991, Bio/Technology 9, pp. 957-962), Maize (Gordon-Kamm et al. 1990, Plant Cell 2, pp. 603-618), Oat (Somers et al. 1992, Bio/Technology 10, pp 1589-1594), Wheat (Vasil et al. 1991, Bio/Technology 10, pp. 667-674, Weeks et al. 1993, Plant Physiology 102, pp. 1077-1084) and barley (Wan and Lemaux 1994, Plant Physiology 102, pp. 37-48, review Vasil 1994, Plant Mol. Biol. 25, pp 925-937).

More specifically, *Agrobacterium* mediated transformation is conveniently achieved as follows:

A vector system carrying the maltogenic alpha-amylase is constructed. The vector system may comprise one vector, but it can comprise two vectors. In the case of two vectors the vector system is referred to as a binary vector system (Gynheung An et al.(1980), Binary Vectors, Plant Molecular Biology Manual A3, 1-19).

An *Agrobacterium* based plant transformation vector consists of replication origin(s) for both *E.coli* and *Agrobacterium* and a bacterial selection marker. A right and preferably also a left border from the Ti plasmid from *Agrobacterium tumefaciens* or from the Ri plasmid from *Agrobacterium rhizogens* is necessary for the transformation of the plant. Between the borders the expression construct is placed which contains the maltogenic alpha-amylase gene and appropriate regulatory sequences such as promotor and terminator sequences. Additionally, a selection gene e.g. the neomycin phosphotransferase type II (NPTII) gene from transposon Tn5 and a reporter gene such as the GUS (beta-glucuronidase) gene is cloned between the borders. A disabled *Agrobacterium* strain harboring a helper plasmid containing the virulens genes is transformed with the above vector. The transformed *Agrobacterium* strain is then used for plant transformation.

### Industrial processes

In principle, the seeds of the invention may be used in any industrial process for which purified maltogenic alpha-amylase or beta-amylase are normally used to catalyze a reaction

between one or more substrate so as to produce the desired effects or products. Of particular interest for the present invention is the use of the seeds in the bread making process for improving the properties of a dough or a baked product.

- 5 According to one embodiment of the present invention the seeds of the invention are used directly in the baking process without the need for first extracting and/or isolating the enzyme. For use in a baking process it is preferred that the seeds containing the maltogenic alpha-amylase or beta-amylase  
10 are milled so as to obtain a consistency suitable for baking.

According to one aspect of the invention the seeds, optionally in a ground form, are used for preparing a flour, in particular wheat flour. More specifically, the flour may be prepared by  
15 milling seeds of the invention containing a maltogenic alpha-amylase or a beta-amylase. The milling may be conducted in accordance with methods known in the art for preparing flour from seeds.

- 20 When a flour has been produced from seeds of the present invention the maltogenic alpha-amylase activity of the resulting flour is normally measured and the strength of the enzyme activity adjusted. For instance, if too much maltogenic alpha-amylase activity is present in the flour prepared from  
25 transgenic seeds of the invention the flour may be diluted with flour free from the maltogenic alpha-amylase. If too little maltogenic alpha-amylase activity is present in the flour additional activity may be added, e.g. in the form of an isolated maltogenic alpha-amylase, such as Novamyl® available  
30 from Novo Nordisk A/S. It follows, that the flour of the present invention may be prepared exclusively from transgenic seeds containing a maltogenic alpha-amylase or from a mixture of seeds which in addition to the transgenic seed of the invention contains non-transgenic seeds or seeds which  
35 otherwise do not contain the maltogenic alpha-amylase. The seeds of the invention preferably contains the maltogenic alpha-amylase in an amount which is effective to delay staling of a baked bread based on said seeds. One embodiment, the seed



of the invention contains a measurable amount of the maltogenic alpha-amylase or the beta-amylase.

The flour of the invention may be is used in accordance with conventional techniques for the production of baked products, in particular bread products. The resulting baked product has an improved anti-staling effect, i.e. the baked product has a reduced rate of deterioration of quality parameters, e.g., softness and/or elasticity, during storage.

The maltogenic alpha-amylase, Novamyl®, has a very unique performance in bread making. Other thermostable  $\alpha$ -amylases like BAN® or Termamyl® must be dosed very carefully in tight intervals, e.g. between 0.5-2 times of the optimum dosage in a given recipe. Otherwise the risk is high that there is either no effect (low dosage) or too high effect (high dosage). The latter will result in a gummy, non-elastic and sticky crumb, unsuited for eating. The maltogenic alpha-amylase as represented by Novamyl® does not have this problem, but can be dosed broadly. For instance, Novamyl® has a positive function on e.g. staling properties from a level of e.g. 200 MANU/kg flour to 5.000 MANU/kg, i.e. a much safer amylase in practical application than other  $\alpha$ -amylases.

The term "baked product" is defined herein as any product prepared from a dough, either of a soft or a crisp character. Examples of baked products, whether of a white, light or dark type, which may be advantageously produced by the present invention are bread (in particular white, whole-meal or rye bread), typically in the form of loaves or rolls, French baguette-type bread, pasta, pita bread, tortillas, tacos, cakes, pancakes, biscuits, cookies, pie crusts, steamed bread, and crisp bread, and the like.

In terms of enzyme activity, the appropriate dosage of the maltogenic alpha-amylase for exerting a desirable improvement of dough and/or baked products, in particular improved anti-

staling properties, will depend on the specific amylase and the amylase substrate in question. The skilled person may determine a suitable enzyme unit dosage on the basis of methods known in the art. Normally, a suitable dosage of the maltogenic alpha-amylase (as present in the flour) is in the range 200 - 5.000 MANU/kg flour.

#### Determination of maltogenic amylase in MANU

One Maltogenic Amylase-Novo Unit (MANU) is the amount of enzyme which under standard will cleave one  $\mu\text{mol}$  maltotriose per minute. The standard conditions are 10 mg/ml maltotriose, 37°C, pH 5.0, 30 minutes reaction time. The pH dependence is found by repeating this measurement at the same conditions, but at different pH values.

The invention is further illustrated with reference to the following examples which are not intended to be in any way limiting to the scope of the invention as claimed.

#### EXAMPLES

##### Example 1

##### Plasmid construction

The plant novamyl plasmid pNP110 is constructed from the plasmid pAHC25(Christensen. A.H. Sharrock, R.A. and Quail, P.H. (1992) Plant Mol. Bio. 1 18 675-689) containing the *Uda* reporter gene encoding beta-glucuronidase (GUS) and the *bar* gene as selective marker encoding phosphinothricin acetyl transferase which inactivates phosphinothricin, the active component in the herbicides Basta and Bialaphos. Each driven by the maize *ub1* promoter and the first intron and terminated by the polyadenylation signal of *nos3'* gene from *Agrobacterium tumefaciens*. The Novamyl mature gene is amplified using the forward primer: FNP110: 5'-tcccccgggatgagcagttccgcaagcgtcaaa-3' and the reverse primer RNP110: 5'-cgatgagctcctagttttgccacgt-3'

using the pDN452 plasmid as template (DIDERICHSEN B. and CHRISTIANSEN L.(1988)FEMS Microbiol. Lett. 56:53-60) under standard PCR conditions. The fragment of 2.0 Kb is digested with SmaI and SacI and ligated with the vector fragment of the plasmid pAHC25 digested with SmaI and SacI. The obtained plasmid designated pNP110 is used for the transformation experiments.

#### Transformation of Wheat

##### 10 Plant material:

Wheat (*Triticum aestivum* L.) plants are grown in greenhouses or in growth chambers in 16h light ( $350 \mu\text{mol m}^{-2}\text{s}^{-1}$ )/8h dark period at 16°C.

Wheat spikes are harvested when embryos are 1-2 mm. Caryopses are removed from the middle half of spikes 12 days after anthesis and surface sterilized for 10 min 5.25% sodium hypochlorite under stirring and finally washed twice in sterile H<sub>2</sub>O. Immature embryos are dissected from caryopses under a stereomicroscope using a scalpel and transferred to Petri dishes containing MS medium with scutellum side up. Twenty immature embryos are placed side by side in an area of 1cm x 1cm and are ready for bombardment the following day.

##### Culture media:

##### 25 Murashige and Skoog medium for immature embryos (MS):

4.3 g MS salts (Sigma M5524)

25 g Sucrose

100 mg Myo-inositol

500 mg Glutamin

##### 30 100 mg Casein hydrolysate

5 g agarose

H<sub>2</sub>O to 1 L and adjust pH to 5.8.

After autoclaving add 1 ml filter sterilized vitamin solution (10 mg thiamin, 50 mg nicotinic acid, 50 mg pyridoxine HCl and

35 200 mg glycine in 100 mL H<sub>2</sub>O) and 1 mg 2,4-dichlorophenoxyacetic acid (2,4-D).

##### Selection medium (MSS):

As MS without glutamine and casein hydrolysate, but with addition of 3 mg/L Bialaphos after autoclaving.

Shoot induction medium MSSI:

5 2.5 g MS salts; 15 g Sucrose; 50 mg Myo-inositol; 2.5 g agarose;

H<sub>2</sub>O to 1 l and adjust pH to 5.8.

After autoclaving add filter-sterilized 5 mg/l bialaphos and 0.5 ml/l vitamin solution as MS and 0.1 mg/l filter-sterilized

10 BAP, 6-benzylaminopurine

Root induction medium

Standard MS-medium (Sigma M9274) with filter-sterilized 1 mg/l bialaphos

15

Gold Coating:

6 mg gold particles are sterilized in 100µl EtOH and vortexed for 3 min. After centrifugation at 10 K for 1 min and washed twice in H<sub>2</sub>O and finally the gold particles are resuspended in 20 100 µl H<sub>2</sub>O.

15 µg pNP110 Plasmid DNA, 50 µl of 2.5 M CaCl<sub>2</sub> and 20 µl of 0.1 M spermidine are mixed with 50 µl gold suspension during vortexing for 3 min and centrifuged at 500 rpm for 5 min at 4°C. Supernatant is removed and the pellet is resuspended in 25 500 µl EtOH and centrifuged 500 rpm for 5 min at 4°C.

Finally, the pellet is resuspended in 80 µl EtOH and 10 µl coated gold particles are transferred to macrocarriers soaked in 70% EtOH for 10 min and air dried.

30 Bombardment of embryos:

The bombardment chamber and the acceleration cylinder is sterilized by spraying with 70% EtOH and air dried. The delivery pressure is set to 1300 psi.

The rupture disk (1100 psi) and the steel mesh are soaked in 35 EtOH for 10 min and air dried. Rupture disk is placed properly in the holder and is fastened tightly. The macrocarrier and the steel mesh in the assembly unit is placed properly in the chamber (level 2 from top). The Petri dish with immature

embryos is placed properly in the chamber (level 4 from top). Vacuum is turn on and subsequently the pressure is turn on. at 1100 psi the rupture dish break and the DNA is bombarded into the immature embryos. The Petri dish is transferred to a growth chamber at 25°C for 20 h without light.

The following day, the bombarded embryos are spread all over the area of the Petri dish containing MSS medium. After two to three transfers, one each second week, selected callus is transferred to shoot induction (MSSI) medium and transferred to a growth chamber with a 16 h light/8 h dark period. After two weeks green areas and shoot formation are visible. Only green-shoot-callus is transferred to new Petri dishes with MSSI medium or to tubes with root induction medium for two more weeks. Plant with roots in tubes are transferred to soil and placed in a greenhouse for three to four months and mature seeds are harvested.

#### Verification of Transgenic wheat

##### Semipurification of genomic DNA

0.25 g plant material in a eppendorf tube is chilled in N<sub>2</sub> and grinded.

500 µl phenol/chloroform 1:1 and 500 µl buffer(50 mM Tris-HCL, pH 9,0+150 mM LiCl, 5 mM EDTA, pH 8.0 + 5% SDS in H<sub>2</sub>O) is added with 10 µl RNase (10 mg/ml).

Centrifuged 15000 rpm for 10 min

Topfase is transferred to a new eppendorf tube and 500 µl chloroform is added. Centrifuged 15000 rpm for 5 min and topfase is transferred to a new eppendorf tube and 1/10<sup>th</sup> vol 3 M NaAc, pH 5.3 is added with 2 vol EtOH. The tube is placed on ice for 30-60 min.

Centrifuged 1500 rpm for 20 min.

Washed with 70% EtOH, spin 6 min and air dry for 15 min.

Resuspended in 30 H<sub>2</sub>O and check 3 µl in an agarose gel.

##### PCR analysis:

To test for the presence of The novamyl gene in genomic DNA of transformed lines, 250 ng genomic DNA of each transgenic line

is used as template in PCR using the forward primer FNP110:  
5'-tcccccgggatgagcagttccgcaagcgtcaaa-3'

and the reverse primer RNP110: 5'-cgatgagctcctagttttgccacgt-3'.

Standard PCR conditions are used with 40 cycles of 1 min at  
5 94°C, 1 min at 61°C, 2 min 72°C.

Novamyl Positive plant lines showed a band of 2.0 kb, whereas  
non transformed plants showed no fragments of 2.0 kb

### Example 2

10 The nucleotide sequence encoding Novamyl (SEQ ID NO: 1) is  
operably linked to the wheat promoter expressing  $\alpha$ -amylase in  
wheat seeds as described in "Promoter and genotype dependent  
transient expression of a reporter gene in plant  
protoplasts."; Stefanov I ; Ilubaev S ; Feher A ; Margoczi K ;  
15 Dudits D; Acta Biologica Hungarica Vol. 42 , No. 4 pp. 323-330  
( 1991 The resulting DNA construct is inserted into a plasmid  
containing suitable regulatory elements and a selection marker,  
such as described in Example 1.

20 Protoplasts are isolated from wheat cell lines as described in  
("Culture of and fertile plant-regeneration from regenerable  
embryogenic suspension cell-derived protoplasts of  
wheat(triticum-aestivum l)" ; Ahmed, KZ ; Sagi, F; PLANT CELL  
REPORTS Vol. 12 , pp. 175-179 (1993).

25 The nucloetide construct containing the maltogenic alpha-  
amylase coding sequence is inserted into wheat protoplast cells  
via PEG treatment as described in "Factors affecting transient  
expression of vector constructs in wheat protoplasts."; Ahmed  
30 KZ ; Omirulleh S ; Sagi F ; Dudits D; Acta Biologica Hungarica  
Vol. 48 , No. 2 pp. 209-220 (1997). The resulting protoplast is  
regenerated into a wheat plant as described in "Fertile wheat  
(Triticum aestivum L.) regenerants from protoplasts of  
embryogenic suspension culture."; Pauk J ; Kertesz Z ; Jenes B  
35 ; Purnhauser L ; Manninen O ; Pulli S ; Barbas Z ; Dudits D;  
Plant Cell Tissue and Organ Culture Vol. 38, No. 1 pp. 1-10  
(1994). The seeds are harvested, and multiplied and used for

producing transgenic wheat plant expressing Novamyl in its seeds.

### Example 3

- 5 The wheat seeds are milled in accordance with conventional techniques for the preparation of wheat flour. Optionally, the wheat is allowed to malt to a predetermined degree before milling. This will allow a greater expression of the bacterial enzyme. The Novamyl content of the flour is determined in MANU:
- 10 One MANU (Maltogenic Amylase Novo Unit) is defined as the amount of enzyme required to release one mmol of maltose per minute at a concentration of 10 mg of maltotriose (Sigma M 8378) substrate per ml of 0.1 M citrate buffer, pH 5.0 at 37 °C for 30 minutes. If needed the content of Novamyl in the flour is adjusted as
- 15 discussed above in the Detailed Description so as to result in a Novamyl content per kg of flour in the range of 200-5000 MANU/kg of flour.

### Example 4

- 20 A baking trial is carried out. The transformed flour is compared to the original un-transformed wheat "sort". The optimum water absorption is determined on a Farinograph (AACC method The Farinograph Handbook, 3rd Edition, 1984, AACC, Edited by Bert L. D'Appolonia and Wallace H. Kunerth, ISBN 0-913250-37-6).
- 25

### Preparation of White Bread (I)

- The straight-dough bread-making method may be used according to AACC Method 10-10B (in Approved Methods of the American
- 30 Association of Cereal Chemists, Ninth Edition, March 1995; AACC, St. Paul MN, USA).

Basic recipe

Wheat flour	100%
Salt	1.5%
Yeast (fresh)	5.3%
5 Sugar	6.0%
Shortening	3.0%
Water	optimum

All percentages are by weight relative to the wheat flour.

Procedure

1. Dough mixing (Hobart mixer):

The mixing time and speed should be determined by the skilled baker so as to obtain an optimum dough consistency under the testing conditions used.

2. 1st punch (e.g., 52 minutes after start)

3. 2nd punch (e.g., 25 minutes later)

4. Molding and panning (e.g., 13 minutes later).

5. Proofing to desired height (e.g., 33 minutes at 32°C, 82% RH)

5. Baking (e.g., at 215°C for 24 minutes)

Preparation of White Bread (II):

The sponge-dough bread-making method may be used according to AACC Method 10-11 (in Approved Methods of the American Association of Cereal Chemists, Ninth Edition, March 1995; AACC, St. Paul MN, USA).

Basic recipe for Sponge

Wheat flour	60%
Yeast (compressed)	36%
Yeast Food	2%
Water	36%

All percentages are by weight relative to the wheat flour.

Procedure

1. Add water to compressed yeast



2. Add yeast food in dry form with flour
3. Mix sponge (Hobart A-120; Hobart Corp., Troy OH, USA):
  - 0.5 minute at 1<sup>st</sup> speed
  - 1 minute at 2<sup>nd</sup> speed

5 The mixing time may be adjusted so as to obtain an optimum dough consistency under the testing conditions used.

4. Ferment in a fermentation cabinet: 4 hours at 30°C, 85% RH

#### Basic recipe for Dough

10	Wheat flour	40%
	Water	24%
	Sugar	5%
	Shortening	3%
	Salt	2%

15 All percentages are by weight relative to the wheat flour.

#### Procedure

1. Add dough ingredients; begin mixer (1<sup>st</sup> speed)
- 20 2. Add sponge in three approximately equal portions at 15, 25, and 35 seconds mixing time; total mixing time: 1 minute
3. At 2<sup>nd</sup> speed, mix to obtain an optimum dough consistency
4. Ferment in a fermentation cabinet: 30 minutes at 30°C, 85% RH
- 25 5. Intermediate proof: 12-15 minutes in fermentation cabinet
6. Mold and final proof at 35.5°C, 92% RH
7. Bake: 25 minutes at 218°C

#### **Example 5**

- 30 Evaluation of Staling Properties of Bread:

The degree of staling is determined on bread, e.g., on day 1, 3, 7 and 9 after baking. Evaluation of staleness and texture can be done according to AACC method 74-09. The principles for determination of softness and elasticity of bread crumb are as follows:

- 35 1. A slice of bread is compressed with a constant speed in a texture analyser, measuring the force for compression in g.

2. The softness of the crumb is measured as the force at 25% compression.

3. The force at 40% compression (P2) and after keeping 40% compression constant for 30 seconds (P3) is measured. The  
5 ratio (P3/P2) is the elasticity of the crumb.

07-12-2000

NOVEL MALTogenic AMYLASE

170 111200007

170 111200007

DK 009900624

5753 WO: New claims filed December 2000 - response 1st. written opinion

ART 34 AMEND

# CLAIMS

1. A transgenic cereal plant cell comprising a nucleotide sequence encoding a maltogenic alpha-amylase which nucleotide sequence, in the cell, is operably linked to elements required for mediating expression of the nucleotide sequence in the seeds of a cereal plant regenerated from said plant cell, said maltogenic alpha-amylase having an amino acid sequence which has:

- i) at least 70% identity to SEQ ID NO: 2, preferably at least 75%, 80%, 85% or at least 90%, e.g. at least 95%, 97%, 98%, or at least 99%; or
- ii) at least 70% identity to the amino acid sequence set forth in amino acids 1-686 of SEQ ID NO:1, preferably at least 75%, 80%, 85% or at least 90%, e.g. at least 95%, 97%, 98%, or at least 99%.

2. The plant cell according to claim 1, wherein the plant cell is a wheat plant cell.

3. The plant cell according to claim 1 or 2, wherein said maltogenic alpha-amylase has the amino acid sequence shown in SEQ ID NO: 2 or the amino sequence acid sequence set forth in amino acids 1-686 of SEQ ID NO:1.

4. The plant cell according to any of the preceding claims, wherein the maltogenic alpha-amylase in baking has an antistaling effect and can be dosed broadly essentially without negative effect.

5. The plant cell according to any of the preceding claims, wherein a seed specific promoter drives the expression of the maltogenic alpha-amylase.

6. The plant cell according to any of the preceding claims, wherein the nucleotide sequence encoding the maltogenic alpha-amylase is derived from a microorganism, preferably a bacterium.

09831656-090201

07-12-2000

5753-WO: New claims filed December 2000 - response 1st. written opinion

DK 009900624

ART 34 AAMD

7. The plant cell according to claim 6, wherein the nucleotide sequence encoding the maltogenic alpha-amylase is derived from the *Bacillus* strain NCIB 11837.

8. A transgenic cereal plant regenerated from a plant cell according to any of the preceding claims and the progeny of said parent plant, which parent or progeny plant is capable of expressing said maltogenic alpha-amylase in the seeds.

9. A transgenic cereal plant comprising a nucleotide sequence encoding a maltogenic alpha-amylase which nucleotide sequence is operably linked to elements required for mediating expression of the nucleotide sequence in the seeds of said plant, wherein said maltogenic alpha-amylase has an amino acid sequence having:

- i) at least 70% identity to SEQ ID NO: 2, preferably at least 75%, 80%, 85% or at least 90%, e.g. at least 95%, 97%, 98%, or at least 99%; or
- ii) at least 70% identity to the amino acid sequence set forth in amino acids 1-686 of SEQ ID NO:1, preferably at least 75%, 80%, 85% or at least 90%, e.g. at least 95%, 97%, 98%, or at least 99%.

10. The plant according to claim 8 or 9 which is wheat.

11. The plant according to any of claims 8-10, wherein said maltogenic amylase is as defined in claims 3 or 4.

12. A seed of a cereal plant defined in any of claims 8-11, said seed containing the maltogenic alpha-amylase in an amount which is effective to delay staling of bread baked from the seed.

13. A transgenic cereal seed comprising maltogenic alpha-amylase in an amount which is effective to delay staling of bread baked from said seed, wherein said maltogenic alpha-amylase has an amino acid sequence having:

07-12-2000

DK 009900624

33-WO: New claims filed December 2000 - response 1st. written opinion

ART 34 AMEND

- i) at least 70% identity to SEQ ID NO: 2, preferably at least 75%, 80%, 85% or at least 90%, e.g. at least 95%, 97%, 98%, or at least 99%; or
- ii) at least 70% identity to the amino acid sequence set forth in amino acids 1-686 of SEQ ID NO:1, preferably at least 75%, 80%, 85% or at least 90%, e.g. at least 95%, 97%, 98%, or at least 99%.

14. The seed of claim 12 or 13, wherein said maltogenic alpha-amylase is as defined in any of claims 3 or 4.

15. The seed of any of claims 12-14, wherein said seed is wheat.

16. Use of seeds as defined in any of claims 12-15 for the preparation of flour or dough.

17. Use of seeds as defined in any of claims 12-15 in baking.

18. A method for preparing a baked product, said method comprising the steps of:

i) expressing a maltogenic alpha-amylase in the seeds of a transgenic cereal plant, said maltogenic alpha-amylase having an amino acid sequence which has:

- a) at least 70% identity to SEQ ID NO: 2, preferably at least 75%, 80%, 85% or at least 90%, e.g. at least 95%, 97%, 98%, or at least 99%; or
- b) at least 70% identity to the amino acid sequence set forth in amino acids 1-686 of SEQ ID NO:1, preferably at least 75%, 80%, 85% or at least 90%, e.g. at least 95%, 97%, 98%, or at least 99%;

ii) preparing flour from said cereal seed comprising said maltogenic alpha-amylase;

iii) preparing a dough comprising the flour of step ii); and

iv) baking the dough to obtain a baked product.

19. A method for preparing a baked product, said method comprising the steps of:

07-12-2000

5753-110: New claims filed December 2000 - response 1st. written opinion

DK 009900624

ART 34 AMEND

i) preparing flour from cereal seed, said seed comprising a maltogenic alpha-amylase having an amino acid sequence which has:

- a) at least 70% identity to SEQ ID NO: 2, preferably at least 75%, 80%, 85% or at least 90%, e.g. at least 95%, 97%, 98%, or at least 99%; or
- b) at least 70% identity to the amino acid sequence set forth in amino acids 1-686 of SEQ ID NO:1, preferably at least 75%, 80%, 85% or at least 90%, e.g. at least 95%, 97%, 98%, or at least 99%;

ii) preparing a dough comprising the flour of step i); and  
iii) baking the dough to obtain a baked product.

20. A method for preparing a baked product, said method comprising the steps of:

i) preparing a dough from flour obtained from cereal seed, said seed comprising a maltogenic alpha-amylase having an amino acid sequence which has:

- a) at least 70% identity to SEQ ID NO: 2, preferably at least 75%, 80%, 85% or at least 90%, e.g. at least 95%, 97%, 98%, or at least 99%; or
- b) at least 70% identity to the amino acid sequence set forth in amino acids 1-686 of SEQ ID NO:1, preferably at least 75%, 80%, 85% or at least 90%, e.g. at least 95%, 97%, 98%, or at least 99%;

ii) preparing a dough comprising the flour of step i); and  
iii) baking the dough to obtain a baked product.

21. The method according to any of claims 18-20, wherein the maltogenic alpha-amylase is as defined in claim 2 or 3.

22. The method according to any of claims 18-20, wherein said seed is as defined in any of claims 12-15.

# SEQUENCE LISTING

<110> Novozymes A/S  
Nielsen, Jack Bech  
Kjaerulff, Soren

<120> Transgenic Plant Expressing Maltogenic Alpha-Amylase

<130> 5753.204-US

<140> 09/831,656

<141> 2001-05-11

<150> PCT/DK99/00624

<151> 1999-11-12

<160> 4

<170> PatentIn version 3.1

<210> 1

<211> 2160

<212> DNA

<213> Bacillus sp.

<400> 1

```
atgaaaaaga aaacgctttc tttatttggt ggactgatgc tctcatcgg tcttctgttc      60
agcggttctc ttccgtacaa tccaaacgcc gctgaagcca gcagttccgc aagcgtcaaa      120
ggggacgtga ttaccagat tatcattgac cggttttacg atggggacac gacgaacaac      180
aatcctgcca aaagttatgg actttacgat ccgaccaaatt cgaagtggaa aatgtattgg      240
ggcgggggatc tggaggggggt tcgtcaaaaa cttccttatc ttaaacagct gggcgtaacg      300
acaatctggt tgtccccggt tttggacaat ctggatacac tggcgggcac cgataacacg      360
ggctatcacg gatactggac gcgcgatttt aaacagattg aggaacattt cgggaattgg      420
accacatttg acacgttggt caatgatgct caccaaaacg gaatcaaggt gattgtcgac      480
tttgtgcccc atcattcgac tccttttaag gcaaacgatt ccacctttgc ggaaggcggc      540
gccctctaca acaatggaac ctatatgggc aattattttg atgacgcaac aaaagggtac      600
ttccaccata atggggacat cagcaactgg gacgaccggt acgaggcgca atggaaaaac      660
ttcacggatc cagccggttt ctcgcttgcc gatttgtcgc aggaaaatgg cagcattgct      720
caatacctga ccgatgcggc ggttcaattg gtagcacatg gagcggatgg tttgcggatt      780
```

gatgcggtga agcatttttaa ttcgggggttc tccaaatcgt tggccgataa actgtaccaa	840
aagaaagaca ttttcctggt gggggaatgg tacggagatg accccggaac agccaatcat	900
ctggaaaagg tccggtacgc caacaacagc ggtgtcaatg tgctggattt tgatctcaac	960
acggtgattc gaaatgtggt cggcacattt acgcaaacga tgtacgatct taacaatatg	1020
gtgaaccaa cggggaacga gtacaaatac aaagaaaatc taatcacatt tatcgataac	1080
catgatatgt caagatttct ttcggtaaat tcgaacaagg cgaatttgca ccaggcgctt	1140
gctttcattc tcaacttcgog gggtagcccc tccatctatt atggaaccga acaatacatg	1200
gcaggcgga atgaccogta caaccggggg atgatgccgg cgtttgatac gacaaccacc	1260
gcctttaaag aggtgtcaac tctggcgggg ttgcgagga acaatgcggc gatccagtac	1320
ggcaccacca cccagcggtg gatcaacaat gatgtttaca tttatgaacg gaaatttttc	1380
aacgatgtcg tgttggtggc catcaatcga aacacgcaat cctcctattc gatttcgggt	1440
ttgcagacgg ccttgccaaa tggcagctat gcggtattatc tgtcagggtt gttggggggg	1500
aacgggattt ccgtttccaa tggaaagtgc gcttcgttca cgtttgcgcc tggagccgtg	1560
tctgtttggc agtacagcac atccgcttca gcgccgcaaa tcggatcggt tgctccaaat	1620
atggggattc cgggtaatgt ggtcacgac gacgggaaag gttttgggac gacgcagga	1680
accgtgacat ttggcggagt gacagcgact gtgaaatcct ggacatccaa tcggattgaa	1740
gtgtacgttc ccaacatggc cgccgggctg accgatgtga aagtcaccgc gggaggagtt	1800
tccagcaatc tgtattctta caatattttg agtggaacgc agacatcggt tgtgtttact	1860
gtgaaaagtg cgcctccgac caacctgggg gataagattt acctgacggg caacataccg	1920
gaattgggga attggagcac ggatacgagc ggagccgtta acaatgcgca agggcccctg	1980
ctcgcgcca attatccgga ttggttttat gtattcagcg ttccagcagg aaagacgatt	2040
caattcaagt tcttcatcaa gcgtgcggat ggaacgattc aatgggagaa tggttcgaac	2100
cacgtggcca caactcccac gggtgcaacc ggtaacatta ctgttacgtg gcaaaactag	2160

<210> 2  
 <211> 719  
 <212> PRT  
 <213> Bacillus sp



<400> 2

Met Lys Lys Lys Thr Leu Ser Leu Phe Val Gly Leu Met Leu Leu Ile  
1 5 10 15

Gly Leu Leu Phe Ser Gly Ser Leu Pro Tyr Asn Pro Asn Ala Ala Glu  
20 25 30

Ala Ser Ser Ser Ala Ser Val Lys Gly Asp Val Ile Tyr Gln Ile Ile  
35 40 45

Ile Asp Arg Phe Tyr Asp Gly Asp Thr Thr Asn Asn Asn Pro Ala Lys  
50 55 60

Ser Tyr Gly Leu Tyr Asp Pro Thr Lys Ser Lys Trp Lys Met Tyr Trp  
65 70 75 80

Gly Gly Asp Leu Glu Gly Val Arg Gln Lys Leu Pro Tyr Leu Lys Gln  
85 90 95

Leu Gly Val Thr Thr Ile Trp Leu Ser Pro Val Leu Asp Asn Leu Asp  
100 105 110

Thr Leu Ala Gly Thr Asp Asn Thr Gly Tyr His Gly Tyr Trp Thr Arg  
115 120 125

Asp Phe Lys Gln Ile Glu Glu His Phe Gly Asn Trp Thr Thr Phe Asp  
130 135 140

Thr Leu Val Asn Asp Ala His Gln Asn Gly Ile Lys Val Ile Val Asp  
145 150 155 160

Phe Val Pro Asn His Ser Thr Pro Phe Lys Ala Asn Asp Ser Thr Phe  
165 170 175

Ala Glu Gly Gly Ala Leu Tyr Asn Asn Gly Thr Tyr Met Gly Asn Tyr  
180 185 190

Phe Asp Asp Ala Thr Lys Gly Tyr Phe His His Asn Gly Asp Ile Ser  
195 200 205

Asn Trp Asp Asp Arg Tyr Glu Ala Gln Trp Lys Asn Phe Thr Asp Pro  
210 215 220

Ala Gly Phe Ser Leu Ala Asp Leu Ser Gln Glu Asn Gly Thr Ile Ala  
225 230 235 240

Gln Tyr Leu Thr Asp Ala Ala Val Gln Leu Val Ala His Gly Ala Asp  
245 250 255

Gly Leu Arg Ile Asp Ala Val Lys His Phe Asn Ser Gly Phe Ser Lys  
260 265 270

Ser Leu Ala Asp Lys Leu Tyr Gln Lys Lys Asp Ile Phe Leu Val Gly  
275 280 285

Glu Trp Tyr Gly Asp Asp Pro Gly Thr Ala Asn His Leu Glu Lys Val  
290 295 300

Arg Tyr Ala Asn Asn Ser Gly Val Asn Val Leu Asp Phe Asp Leu Asn  
305 310 315 320

Thr Val Ile Arg Asn Val Phe Gly Thr Phe Thr Gln Thr Met Tyr Asp  
325 330 335

Leu Asn Asn Met Val Asn Gln Thr Gly Asn Glu Tyr Lys Tyr Lys Glu  
340 345 350

Asn Leu Ile Thr Phe Ile Asp Asn His Asp Met Ser Arg Phe Leu Ser  
355 360 365

Val Asn Ser Asn Lys Ala Asn Leu His Gln Ala Leu Ala Phe Ile Leu  
370 375 380

Thr Ser Arg Gly Thr Pro Ser Ile Tyr Tyr Gly Thr Glu Gln Tyr Met  
385 390 395 400

Ala Gly Gly Asn Asp Pro Tyr Asn Arg Gly Met Met Pro Ala Phe Asp  
405 410 415



Protein Data Bank

Pro Pro Thr Asn Leu Gly Asp Lys Ile Tyr Leu Thr Gly Asn Ile Pro  
625 630 635 640

Glu Leu Gly Asn Trp Ser Thr Asp Thr Ser Gly Ala Val Asn Asn Ala  
645 650 655

Gln Gly Pro Leu Leu Ala Pro Asn Tyr Pro Asp Trp Phe Tyr Val Phe  
660 665 670

Ser Val Pro Ala Gly Lys Thr Ile Gln Phe Lys Phe Phe Ile Lys Arg  
675 680 685

Ala Asp Gly Thr Ile Gln Trp Glu Asn Gly Ser Asn His Val Ala Thr  
690 695 700

Thr Pro Thr Gly Ala Thr Gly Asn Ile Thr Val Thr Trp Gln Asn  
705 710 715

<210> 3  
<211> 33  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> FNP 110

<400> 3  
tccccgggga tgagcagttc cgcaagcgtc aaa 33

<210> 4  
<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> RNP 110

<400> 4  
cgatgagctc ctagttttgc cacgt 25

## SEQUENCE LISTING

&lt;110&gt; Novo Nordisk A/S

&lt;120&gt; TRANSGENIC PLANT EXPRESSING MALTOGENIC ALPHA-AMYLASE

&lt;130&gt; 5753A

&lt;140&gt;

&lt;141&gt;

&lt;160&gt; 4

&lt;170&gt; PatentIn Ver. 2.1

&lt;210&gt; 1

&lt;211&gt; 2160

&lt;212&gt; DNA

&lt;213&gt; Bacillus sp.

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)..(2160)

&lt;220&gt;

&lt;221&gt; mat\_peptide

&lt;222&gt; (100)..(2157)

&lt;400&gt; 1

atg	aaa	aag	aaa	acg	ctt	tct	tta	ttt	gtg	gga	ctg	atg	ctc	ctc	atc	48
Met	Lys	Lys	Lys	Thr	Leu	Ser	Leu	Phe	Val	Gly	Leu	Met	Leu	Leu	Ile	
				-30				-25							-20	

ggt	ctt	ctg	ttc	agc	ggt	tct	ctt	ccg	tac	aat	cca	aac	gcc	gct	gaa	96
Gly	Leu	Leu	Phe	Ser	Gly	Ser	Leu	Pro	Tyr	Asn	Pro	Asn	Ala	Ala	Glu	
			-15				-10								-5	

gcc	agc	agt	tcc	gca	agc	gtc	aaa	ggg	gac	gtg	att	tac	cag	att	atc	144
Ala	Ser	Ser	Ser	Ala	Ser	Val	Lys	Gly	Asp	Val	Ile	Tyr	Gln	Ile	Ile	
-1	1				5				10						15	

att	gac	cgg	ttt	tac	gat	ggg	gac	acg	acg	aac	aac	aat	cct	gcc	aaa	192
Ile	Asp	Arg	Phe	Tyr	Asp	Gly	Asp	Thr	Thr	Asn	Asn	Asn	Pro	Ala	Lys	
				20				25							30	

agt	tat	gga	ctt	tac	gat	ccg	acc	aaa	tcg	aag	tgg	aaa	atg	tat	tgg	240
Ser	Tyr	Gly	Leu	Tyr	Asp	Pro	Thr	Lys	Ser	Lys	Trp	Lys	Met	Tyr	Trp	
			35					40							45	

ggc ggg gat ctg gag ggg gtt cgt caa aaa ctt cct tat ctt aaa cag 288  
 Gly Gly Asp Leu Glu Gly Val Arg Gln Lys Leu Pro Tyr Leu Lys Gln  
 50 55 60

ctg ggc gta acg aca atc tgg ttg tcc ccg gtt ttg gac aat ctg gat 336  
 Leu Gly Val Thr Thr Ile Trp Leu Ser Pro Val Leu Asp Asn Leu Asp  
 65 70 75

aca ctg gcg ggc acc gat aac acg ggc tat cac gga tac tgg acg cgc 384  
 Thr Leu Ala Gly Thr Asp Asn Thr Gly Tyr His Gly Tyr Trp Thr Arg  
 80 85 90 95

gat ttt aaa cag att gag gaa cat ttc ggg aat tgg acc aca ttt gac 432  
 Asp Phe Lys Gln Ile Glu Glu His Phe Gly Asn Trp Thr Thr Phe Asp  
 100 105 110

acg ttg gtc aat gat gct cac caa aac gga atc aag gtg att gtc gac 480  
 Thr Leu Val Asn Asp Ala His Gln Asn Gly Ile Lys Val Ile Val Asp  
 115 120 125

ttt gtg ccc aat cat tcg act cct ttt aag gca aac gat tcc acc ttt 528  
 Phe Val Pro Asn His Ser Thr Pro Phe Lys Ala Asn Asp Ser Thr Phe  
 130 135 140

gcg gaa ggc ggc gcc ctc tac aac aat gga acc tat atg ggc aat tat 576  
 Ala Glu Gly Gly Ala Leu Tyr Asn Asn Gly Thr Tyr Met Gly Asn Tyr  
 145 150 155

ttt gat gac gca aca aaa ggg tac ttc cac cat aat ggg gac atc agc 624  
 Phe Asp Asp Ala Thr Lys Gly Tyr Phe His His Asn Gly Asp Ile Ser  
 160 165 170 175

aac tgg gac gac cgg tac gag gcg caa tgg aaa aac ttc acg gat cca 672  
 Asn Trp Asp Asp Arg Tyr Glu Ala Gln Trp Lys Asn Phe Thr Asp Pro  
 180 185 190

gcc ggt ttc tcg ctt gcc gat ttg tcg cag gaa aat ggc acg att gct 720  
 Ala Gly Phe Ser Leu Ala Asp Leu Ser Gln Glu Asn Gly Thr Ile Ala  
 195 200 205

caa tac ctg acc gat gcg gcg gtt caa ttg gta gca cat gga gcg gat 768  
 Gln Tyr Leu Thr Asp Ala Ala Val Gln Leu Val Ala His Gly Ala Asp  
 210 215 220

ggt ttg cgg att gat gcg gtg aag cat ttt aat tcg ggg ttc tcc aaa 816  
 Gly Leu Arg Ile Asp Ala Val Lys His Phe Asn Ser Gly Phe Ser Lys  
 225 230 235

tcg ttg gcc gat aaa ctg tac caa aag aaa gac att ttc ctg gtg ggg 864  
 Ser Leu Ala Asp Lys Leu Tyr Gln Lys Lys Asp Ile Phe Leu Val Gly  
 240 245 250 255

gaa tgg tac gga gat gac ccc gga aca gcc aat cat ctg gaa aag gtc 912  
 Glu Trp Tyr Gly Asp Asp Pro Gly Thr Ala Asn His Leu Glu Lys Val  
 260 265 270

cgg tac gcc aac aac agc ggt gtc aat gtg ctg gat ttt gat ctc aac 960  
 Arg Tyr Ala Asn Asn Ser Gly Val Asn Val Leu Asp Phe Asp Leu Asn  
 275 280 285

acg gtg att cga aat gtg ttc ggc aca ttt acg caa acg atg tac gat 1008  
 Thr Val Ile Arg Asn Val Phe Gly Thr Phe Thr Gln Thr Met Tyr Asp  
 290 295 300

ctt aac aat atg gtg aac caa acg ggg aac gag tac aaa tac aaa gaa 1056  
 Leu Asn Asn Met Val Asn Gln Thr Gly Asn Glu Tyr Lys Tyr Lys Glu  
 305 310 315

aat cta atc aca ttt atc gat aac cat gat atg tca aga ttt ctt tcg 1104  
 Asn Leu Ile Thr Phe Ile Asp Asn His Asp Met Ser Arg Phe Leu Ser  
 320 325 330 335

gta aat tcg aac aag gcg aat ttg cac cag gcg ctt gct ttc att ctc 1152  
 Val Asn Ser Asn Lys Ala Asn Leu His Gln Ala Leu Ala Phe Ile Leu  
 340 345 350

act tcg cgg ggt acg ccc tcc atc tat tat gga acc gaa caa tac atg 1200  
 Thr Ser Arg Gly Thr Pro Ser Ile Tyr Tyr Gly Thr Glu Gln Tyr Met  
 355 360 365

gca ggc ggc aat gac ccg tac aac cgg ggg atg atg ccg gcg ttt gat 1248  
 Ala Gly Gly Asn Asp Pro Tyr Asn Arg Gly Met Met Pro Ala Phe Asp  
 370 375 380

acg aca acc acc gcc ttt aaa gag gtg tca act ctg gcg ggg ttg cgc 1296  
 Thr Thr Thr Thr Ala Phe Lys Glu Val Ser Thr Leu Ala Gly Leu Arg  
 385 390 395

agg aac aat gcg gcg atc cag tac ggc acc acc acc cag cgt tgg atc 1344  
 Arg Asn Asn Ala Ala Ile Gln Tyr Gly Thr Thr Thr Gln Arg Trp Ile  
 400 405 410 415

aac aat gat gtt tac att tat gaa cgg aaa ttt ttc aac gat gtc gtg 1392  
 Asn Asn Asp Val Tyr Ile Tyr Glu Arg Lys Phe Phe Asn Asp Val Val  
 420 425 430

ttg gtg gcc atc aat cga aac acg caa tcc tcc tat tcg att tcc ggt 1440  
 Leu Val Ala Ile Asn Arg Asn Thr Gln Ser Ser Tyr Ser Ile Ser Gly  
 435 440 445

ttg cag acg gcc ttg cca aat ggc agc tat gcg gat tat ctg tca ggg 1488  
 Leu Gln Thr Ala Leu Pro Asn Gly Ser Tyr Ala Asp Tyr Leu Ser Gly  
 450 455 460

ctg ttg ggg ggg aac ggg att tcc gtt tcc aat gga agt gtc gct tcg 1536  
 Leu Leu Gly Gly Asn Gly Ile Ser Val Ser Asn Gly Ser Val Ala Ser  
 465 470 475

ttc acg ctt gcg cct gga gcc gtg tct gtt tgg cag tac agc aca tcc 1584  
 Phe Thr Leu Ala Pro Gly Ala Val Ser Val Trp Gln Tyr Ser Thr Ser  
 480 485 490 495

gct tca gcg ccg caa atc gga tcg gtt gct cca aat atg ggg att ccg 1632  
 Ala Ser Ala Pro Gln Ile Gly Ser Val Ala Pro Asn Met Gly Ile Pro  
 500 505 510

ggt aat gtg gtc acg atc gac ggg aaa ggt ttt ggg acg acg cag gga 1680  
 Gly Asn Val Val Thr Ile Asp Gly Lys Gly Phe Gly Thr Thr Gln Gly  
 515 520 525

acc gtg aca ttt ggc gga gtg aca gcg act gtg aaa tcc tgg aca tcc 1728  
 Thr Val Thr Phe Gly Gly Val Thr Ala Thr Val Lys Ser Trp Thr Ser  
 530 535 540

aat cgg att gaa gtg tac gtt ccc aac atg gcc gcc ggg ctg acc gat 1776  
 Asn Arg Ile Glu Val Tyr Val Pro Asn Met Ala Ala Gly Leu Thr Asp  
 545 550 555

gtg aaa gtc acc gcg ggt gga gtt tcc agc aat ctg tat tct tac aat 1824  
 Val Lys Val Thr Ala Gly Gly Val Ser Ser Asn Leu Tyr Ser Tyr Asn  
 560 565 570 575

att ttg agt gga acg cag aca tcg gtt gtg ttt act gtg aaa agt gcg 1872  
 Ile Leu Ser Gly Thr Gln Thr Ser Val Val Phe Thr Val Lys Ser Ala  
 580 585 590

cct ccg acc aac ctg ggg gat aag att tac ctg acg ggc aac ata ccg 1920  
 Pro Pro Thr Asn Leu Gly Asp Lys Ile Tyr Leu Thr Gly Asn Ile Pro  
 595 600 605

gaa ttg ggg aat tgg agc acg gat acg agc gga gcc gtt aac aat gcg 1968  
 Glu Leu Gly Asn Trp Ser Thr Asp Thr Ser Gly Ala Val Asn Asn Ala  
 610 615 620



caa ggg ccc ctg ctc gcg ccc aat tat ccg gat tgg ttt tat gta ttc 2016  
 Gln Gly Pro Leu Leu Ala Pro Asn Tyr Pro Asp Trp Phe Tyr Val Phe  
 625 630 635

agc gtt cca gca gga aag acg att caa ttc aag ttc ttc atc aag cgt 2064  
 Ser Val Pro Ala Gly Lys Thr Ile Gln Phe Lys Phe Phe Ile Lys Arg  
 640 645 650 655

gcg gat gga acg att caa tgg gag aat ggt tcg aac cac gtg gcc aca 2112  
 Ala Asp Gly Thr Ile Gln Trp Glu Asn Gly Ser Asn His Val Ala Thr  
 660 665 670

act ccc acg ggt gca acc ggt aac att act gtt acg tgg caa aac tag 2160  
 Thr Pro Thr Gly Ala Thr Gly Asn Ile Thr Val Thr Trp Gln Asn  
 675 680 685

<210> 2

<211> 719

<212> PRT

<213> Bacillus sp.

<400> 2

Met Lys Lys Lys Thr Leu Ser Leu Phe Val Gly Leu Met Leu Leu Ile  
 1 5 10 15

Gly Leu Leu Phe Ser Gly Ser Leu Pro Tyr Asn Pro Asn Ala Ala Glu  
 20 25 30

Ala Ser Ser Ser Ala Ser Val Lys Gly Asp Val Ile Tyr Gln Ile Ile  
 35 40 45

Ile Asp Arg Phe Tyr Asp Gly Asp Thr Thr Asn Asn Asn Pro Ala Lys  
 50 55 60

Ser Tyr Gly Leu Tyr Asp Pro Thr Lys Ser Lys Trp Lys Met Tyr Trp  
 65 70 75 80

Gly Gly Asp Leu Glu Gly Val Arg Gln Lys Leu Pro Tyr Leu Lys Gln  
 85 90 95

Leu Gly Val Thr Thr Ile Trp Leu Ser Pro Val Leu Asp Asn Leu Asp  
 100 105 110

Thr Leu Ala Gly Thr Asp Asn Thr Gly Tyr His Gly Tyr Trp Thr Arg  
 115 120 125

Asp Phe Lys Gln Ile Glu Glu His Phe Gly Asn Trp Thr Thr Phe Asp  
 130 135 140

Thr Leu Val Asn Asp Ala His Gln Asn Gly Ile Lys Val Ile Val Asp  
 145 150 155 160

Phe Val Pro Asn His Ser Thr Pro Phe Lys Ala Asn Asp Ser Thr Phe  
 165 170 175

Ala Glu Gly Gly Ala Leu Tyr Asn Asn Gly Thr Tyr Met Gly Asn Tyr  
 180 185 190

Phe Asp Asp Ala Thr Lys Gly Tyr Phe His His Asn Gly Asp Ile Ser  
 195 200 205

Asn Trp Asp Asp Arg Tyr Glu Ala Gln Trp Lys Asn Phe Thr Asp Pro  
 210 215 220

Ala Gly Phe Ser Leu Ala Asp Leu Ser Gln Glu Asn Gly Thr Ile Ala  
 225 230 235 240

Gln Tyr Leu Thr Asp Ala Ala Val Gln Leu Val Ala His Gly Ala Asp  
 245 250 255

Gly Leu Arg Ile Asp Ala Val Lys His Phe Asn Ser Gly Phe Ser Lys  
 260 265 270

Ser Leu Ala Asp Lys Leu Tyr Gln Lys Lys Asp Ile Phe Leu Val Gly  
 275 280 285

Glu Trp Tyr Gly Asp Asp Pro Gly Thr Ala Asn His Leu Glu Lys Val  
 290 295 300

Arg Tyr Ala Asn Asn Ser Gly Val Asn Val Leu Asp Phe Asp Leu Asn  
 305 310 315 320

Thr Val Ile Arg Asn Val Phe Gly Thr Phe Thr Gln Thr Met Tyr Asp  
 325 330 335

Leu Asn Asn Met Val Asn Gln Thr Gly Asn Glu Tyr Lys Tyr Lys Glu  
 340 345 350

Asn Leu Ile Thr Phe Ile Asp Asn His Asp Met Ser Arg Phe Leu Ser  
 355 360 365

Val Asn Ser Asn Lys Ala Asn Leu His Gln Ala Leu Ala Phe Ile Leu  
 370 375 380



Glu Leu Gly Asn Trp Ser Thr Asp Thr Ser Gly Ala Val Asn Asn Ala  
                     645                    650                    655

Gln Gly Pro Leu Leu Ala Pro Asn Tyr Pro Asp Trp Phe Tyr Val Phe  
                     660                    665                    670

Ser Val Pro Ala Gly Lys Thr Ile Gln Phe Lys Phe Phe Ile Lys Arg  
                     675                    680                    685

Ala Asp Gly Thr Ile Gln Trp Glu Asn Gly Ser Asn His Val Ala Thr  
                     690                    695                    700

Thr Pro Thr Gly Ala Thr Gly Asn Ile Thr Val Thr Trp Gln Asn  
                     705                    710                    715

<210> 3

<211> 33

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:FNP110

<400> 3

tcccccgga tgagcagttc cgcaagcgtc aaa

33

<210> 4

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:RNP110

<400> 4

cgatgagctc ctagttttgc cacgt

25




FIG. 1

Attorney Docket No.: 5753.204-US

PATENT

**COMBINED DECLARATION FOR UTILITY PATENT APPLICATION  
AND POWER OF ATTORNEY**

**As a below named inventor, I hereby declare that:**

My residence, mailing address, and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

**TRANSGENIC PLANT EXPRESSING MALTOGENIC ALPHA-AMYLASE**

(Title of the Invention)

the specification of which

☐ is attached hereto

OR

☒ was filed on (11/12/1999 as PCT International Application Number PCT/DK99/00624 and was amended on (MM/DD/YYYY) \_\_\_\_\_ (if applicable).

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by an amendment referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR 1.56, including for continuation-in-part applications, material information which became available between the filing date of the prior application and the national or PCT international filing date of the continuation-in-part application.

I hereby claim foreign priority benefits under 35 U.S.C. 119(a)-(d) or 365(b) of any foreign application(s) for patent or inventor's certificate, or 365(a) of any PCT international applications(s) for patent or inventor's certificate or of any PCT international applications(s) which designated at least one country other than the United States of America, listed below and have also identified below any foreign application(s) for patent or inventor's certificate, or any PCT international application(s) having a filing date before that of the application(s) on which priority is claimed:

Prior Foreign Application Number(s)	Country	Foreign Filing Date (MM/DD/YYYY)	Priority Claimed
PA 1998 01478	Denmark	11/12/1998	<input checked="" type="checkbox"/> YES <input type="checkbox"/> NO <input type="checkbox"/> YES <input type="checkbox"/> NO <input type="checkbox"/> YES <input type="checkbox"/> NO <input type="checkbox"/> YES <input type="checkbox"/> NO

I hereby claim the benefit under 35 U.S.C. 119(e) of any United States provisional application(s) listed below.

Application Number(s)	Filing Date
60/123,643	03/10/1999



Direct all correspondence to:

25908

PATENT TRADEMARK OFFICE

Customer Number **25908**

I hereby appoint the practitioners at the above mentioned customer number as my/our attorney(s) or agent (s) to prosecute the application identified above, and to transact all business in the United States Patent and Trademark Office connected therewith.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C. 1001 and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

NAME OF SOLE OR FIRST INVENTOR	
Given Name (first and middle [if any])	Jack Bech
Family Name or Surname	Nielsen
Inventor's Signature	<i>Jack B. Nielsen</i>
Date	17/04/2001
Residence: City, State, Country	Hellerup
Citizenship	Danish
Mailing Address	Ole Olsens Allé 12
City, State, Zip, Country	DK-2900 Hellerup, Denmark

NAME OF SECOND INVENTOR	
Given Name (first and middle [if any])	Søren
Family Name or Surname	Kjærulff
Inventor's Signature	<i>Søren Kjærulff</i>
Date	19/4 - 2001
Residence: City, State, Country	Vanløse
Citizenship	Danish
Mailing Address	Kongsdalsvej 47
City, State, Zip, Country	DK-2720 Vanløse, Denmark